

How do normality tests behave for rounded data?

Marina Iturrate–Bobes, Ignacio Montes and Raúl Pérez–Fernández

(iturratemarina, imontes, perezfernandez)@uniovi.es

University of Oviedo



Universidad de Oviedo

ISIPTA'2025, Bielefeld

Who are we?

How do normality
tests behave for
rounded data?

Introduction

Objective

Method



M. Iturrate-Bobes



I. Montes



R. Pérez-Fernández

How do normality
tests behave for
rounded data?

Introduction

Objective

Method




D. Nieto-Barba



M. Iturrate-Bobes



J. Salamanca



I. Montes



E. Miranda



R. Pérez-Fernández

Objective

To study and compare the behaviour of three normality tests in the presence of imprecision caused by rounding:

- 1 Anderson-Darling normality test.
- 2 Lilliefors normality test.
- 3 Shapiro-Wilk normality test.

$$\begin{cases} H_0 : X \equiv N(\mu, \sigma) \\ H_1 : X \not\equiv N(\mu, \sigma) \end{cases}$$

A random set-based approach

- A continuous random variable X is rounded to the d -th decimal number ($d \in \mathbb{Z}$).
- Set of possible values: $\mathbb{Z}_d = \{x \mid 10^d \cdot x \in \mathbb{Z}\}$.
- We construct the random (interval) set:

$$\Gamma = [\tilde{X} - 5 \cdot 10^{-(d+1)}, \tilde{X} + 5 \cdot 10^{-(d+1)}[,$$

where $\tilde{X}$ is the observed rounded variable.

The p-value associated with an interval-valued sample $I_1, \dots, I_n$ is a set

$$\text{p-value}(I_1, \dots, I_n) = \{\text{p-value}(x_1, \dots, x_n) \mid x_i \in I_i, \forall i \in \{1, \dots, n\}\},$$

based on which three regions arise:

(1) rejection region, **(2)** acceptance region and **(3) indecision region**.

Experimental setup

We use Monte Carlo simulation with the aim of:

- 1 Exploring the influence of $\sigma = \sqrt{\text{Var}(X)}$ on the power of the normality tests.
- 2 Determining the probability of the rejection region, the acceptance region and the indecision region for each normality test.

Results can be found in POSTER 6.

Funding from grant PID2022-140585NB-I00 funded by
MICIU/AEI/10.13039/501100011033 and “FEDER/UE”.



How do normality tests behave for rounded data?

Marina Iturrate–Bobes, Ignacio Montes and Raúl Pérez–Fernández

(iturratemarina, imontes, perezfernandez)@uniovi.es

University of Oviedo



Universidad de Oviedo

ISIPTA'2025, Bielefeld