

On the closure of aggregation rules for imprecise probabilities

Enrique Miranda, Ignacio Montes

(mirandaenrique,imontes)@uniovi.es

University of Oviedo



ISIPTA'2025, Bielefeld

Who are we?


Introduction

Aggregation
results

Conclusions



E. Miranda



I. Montes

Who are we?

Introduction

Aggregation
results

Conclusions




M.Iturrate



D.Nieto



E. Miranda



I.Montes



R.Pérez



J.Salamanca

Let $\mathcal{X}$ be a finite space. Assume we have a group of n experts, and that each of them models her uncertainty about an experiment in terms of a coherent lower prevision $\underline{P}_i$ on $\mathcal{L}(\mathcal{X})$, with associated credal set $\mathcal{M}(\underline{P}_i)$.

An **aggregation rule on coherent lower previsions** is a map $\mathcal{A}$ that transforms this input into a lower prevision $\underline{P} := \mathcal{A}(\underline{P}_1, \dots, \underline{P}_n)$ that summarises the opinions of the group.

We consider four aggregation rules and six different families, and study:

- (a) if the aggregation rule is closed for a given family;
- (b) if not, if there are particular cases when it is closed.

Let us meet our contestants

Introduction

Aggregation
results

Conclusions

Conjunction ($\mathcal{A}_C$) It is the lower envelope of $\cap_{i=1}^n \mathcal{M}(\underline{P}_i)$, or, equivalently, the natural extension of $\max \{ \underline{P}_1, \dots, \underline{P}_n \}$.

Let us meet our contestants

Introduction

Aggregation
results

Conclusions

Conjunction ($\mathcal{A}_C$) It is the lower envelope of $\cap_{i=1}^n \mathcal{M}(\underline{P}_i)$, or, equivalently, the natural extension of $\max \{ \underline{P}_1, \dots, \underline{P}_n \}$.

Disjunction ($\mathcal{A}_D$) $\mathcal{A}_D(f) = \min \{ \underline{P}_1(f), \dots, \underline{P}_n(f) \} \forall f$.

Let us meet our contestants

Introduction

Aggregation
results

Conclusions

Conjunction ($\mathcal{A}_C$) It is the lower envelope of $\cap_{i=1}^n \mathcal{M}(\underline{P}_i)$, or, equivalently, the natural extension of $\max \{ \underline{P}_1, \dots, \underline{P}_n \}$.

Disjunction ($\mathcal{A}_D$) $\mathcal{A}_D(f) = \min \{ \underline{P}_1(f), \dots, \underline{P}_n(f) \} \forall f$.

Mixture ($\mathcal{A}_M$) Given $\alpha_1, \dots, \alpha_n \geq 0$ with $\sum_{i=1}^n \alpha_i = 1$, $\mathcal{A}_M(f) = \sum_{j=1}^n \alpha_j \underline{P}_j(f) \forall f$.

Let us meet our contestants

Introduction

Aggregation
results

Conclusions

Conjunction ($\mathcal{A}_C$) It is the lower envelope of $\cap_{i=1}^n \mathcal{M}(\underline{P}_i)$, or, equivalently, the natural extension of $\max \{ \underline{P}_1, \dots, \underline{P}_n \}$.

Disjunction ($\mathcal{A}_D$) $\mathcal{A}_D(f) = \min \{ \underline{P}_1(f), \dots, \underline{P}_n(f) \} \forall f$.

Mixture ($\mathcal{A}_M$) Given $\alpha_1, \dots, \alpha_n \geq 0$ with $\sum_{i=1}^n \alpha_i = 1$, $\mathcal{A}_M(f) = \sum_{j=1}^n \alpha_j \underline{P}_j(f) \forall f$.

Pareto ($\mathcal{A}_P$) $\mathcal{A}_P(f) = \min \{ \max_{j=1, \dots, n} \underline{P}_j(f), \min_{j=1, \dots, n} \overline{P}_j(f) \} \forall f$.

Life is a war

Introduction

Aggregation
results

Conclusions



Comparative probabilities

A **comparative probability model** on singletons is determined by some $\mathcal{L} \subseteq \mathcal{X} \times \mathcal{X}$, so that $\mathcal{M}(\mathcal{L}) = \{P : P(\{x_i\}) \geq P(\{x_j\}) \ \forall (x_i, x_j) \in \mathcal{L}\}$.

- ▶ $\mathcal{A}_C$ is closed for comparative probabilities. ✓✓
- ▶ $\mathcal{A}_D$ is not closed for comparative probabilities, but $\mathcal{A}_D(\underline{P}_1, \underline{P}_2)$ has a unique undominated outer approximation. ✓
- ▶ $\mathcal{A}_M(\underline{P}_1, \underline{P}_2)$ is a comparative probability iff $\mathcal{L}_1 = \mathcal{L}_2$. ✓
- ▶ $\mathcal{A}_P$ is closed for comparative probabilities. ✓✓

2-monotone capacities

Introduction

Aggregation
results

Conclusions

$\underline{P}$ is **2-monotone** when $\underline{P}(A \cup B) + \underline{P}(A \cap B) \geq \underline{P}(A) + \underline{P}(B)$ for any $A, B \subseteq \mathcal{X}$.



2-monotone capacities

$\underline{P}$ is **2-monotone** when $\underline{P}(A \cup B) + \underline{P}(A \cap B) \geq \underline{P}(A) + \underline{P}(B)$ for any $A, B \subseteq \mathcal{X}$.

- ▶ $\mathcal{A}_C$ does not preserve 2-monotonicity, but it does so when $\mathcal{M}(\underline{P}_1) \cup \mathcal{M}(\underline{P}_2)$ is convex. ✓
- ▶ $\mathcal{A}_D$ does not preserve 2-monotonicity, but it does when either $\underline{P}_1$ or $\underline{P}_2$ is a categorical belief function. ✓
- ▶ $\mathcal{A}_M$ preserves 2-monotonicity. ✓✓
- ▶ $\mathcal{A}_P$ does not preserve 2-monotonicity. ✗

Probability intervals

A lower probability $\underline{P}$ is a **probability interval** when $\mathcal{M}(\underline{P})$ is determined by the lower and upper probabilities of the singletons.



Probability intervals

A lower probability $\underline{P}$ is a **probability interval** when $\mathcal{M}(\underline{P})$ is determined by the lower and upper probabilities of the singletons.

- ▶ (De Campos et al. 1994) $\mathcal{A}_C$ preserves being a probability interval. ✓✓
- ▶ (De Campos et al. 1994) $\mathcal{A}_D$ does not preserve being a probability interval, but there is a unique outer approximation of $\mathcal{A}_D(\underline{P}_1, \underline{P}_2)$. ✓
- ▶ $\mathcal{A}_M$ does not preserve being a probability interval, but there is a unique outer approximation of $\mathcal{A}_M(\underline{P}_1, \underline{P}_2)$. ✓
- ▶ $\mathcal{A}_P$ does not preserve being a probability interval, but it does when $\max\{\underline{P}_1, \underline{P}_2\}$ is coherent. ✓

Belief functions

Introduction

Aggregation
results

Conclusions


$\underline{P}$ is a **belief function** (=completely monotone) when its möbius inverse is non-negative.



$\underline{P}$ is a **belief function** (=completely monotone) when its möbius inverse is non-negative.

- ▶ $\mathcal{A}_C$ does not preserve complete monotonicity. ✗
- ▶ $\mathcal{A}_D$ does not preserve complete monotonicity, but it does when either $\underline{P}_1$ or $\underline{P}_2$ is a degenerate probability measure. ✓
- ▶ $\mathcal{A}_M$ preserves complete monotonicity. ✓✓
- ▶ $\mathcal{A}_P$ does not preserve complete monotonicity. ✗

A lower probability $\underline{P}$ is a p -box when $\mathcal{M}(\underline{P})$ is uniquely determined by the lower and upper probabilities of the cumulative events.



A lower probability $\underline{P}$ is a p -box when $\mathcal{M}(\underline{P})$ is uniquely determined by the lower and upper probabilities of the cumulative events.

- ▶ The conjunction $\mathcal{A}_C$ of two p -boxes is a p -box iff $\mathcal{M}(\underline{P}_1) \cap \mathcal{M}(\underline{P}_2) \neq \emptyset$. ✓✓
- ▶ $\mathcal{A}_D$ is not closed in the family of p -boxes, but $\mathcal{A}_D(\underline{P}_1, \underline{P}_2)$ has a unique undominated outer approximation in the family. ✓
- ▶ $\mathcal{A}_M(\underline{P}_1, \underline{P}_2)$ is a p -box iff $\mathcal{F}_1 \cup \mathcal{F}_2$ is totally ordered by set inclusion, where $\mathcal{F}_1, \mathcal{F}_2$ denote the focal elements of $\underline{P}_1, \underline{P}_2$. ✓
- ▶ $\mathcal{A}_P$ is not closed in the family of p -boxes. ✗

Minitive measures

Introduction

Aggregation
results

Conclusions

A lower probability is **minitive** iff $\underline{P}(A \cap B) = \min\{\underline{P}(A), \underline{P}(B)\} \forall A, B \subseteq \mathcal{X}$. Its conjugate $\overline{P}$ is a **maxitive** measure.



Minitive measures

Introduction

Aggregation
results

Conclusions

A lower probability is **minitive** iff $\underline{P}(A \cap B) = \min\{\underline{P}(A), \underline{P}(B)\} \forall A, B \subseteq \mathcal{X}$. Its conjugate $\overline{P}$ is a **maxitive** measure.

- ▶ (Miranda et al. 2015) $\mathcal{A}_C(\overline{P}_1, \overline{P}_2)$ is maxitive iff $\min_i \max_j \overline{P}_i(\{x_j\}) = \max_{j=1,2} \min_{i=1,2} \overline{P}_i(\{x_j\})$ for all $\{x_1, x_2\} \subseteq \mathcal{X}$. ✓
- ▶ $\mathcal{A}_D$ is closed in the family of minitive measures. ✓✓
- ▶ $\mathcal{A}_M(\underline{P}_1, \underline{P}_2)$ is minitive iff $\mathcal{F}_1 \cup \mathcal{F}_2$ is closed under set inclusion. ✓
- ▶ $\mathcal{A}_P$ is not closed in the family of minitive measures. ✗

Summary and conclusions

Introduction

Aggregation
results

Conclusions

	Conj. $\mathcal{A}_C$	Disj. $\mathcal{A}_D$	Mixt. $\mathcal{A}_M$	Pareto $\mathcal{A}_P$
Comparative	✓✓	✓	✓	✓✓
2-monotone	✓	✓	✓✓	✗
Prob. intervals	✓✓	✓	✓	✓
Belief functions	✗	✓	✓✓	✗
p -boxes	✓✓	✓	✓	✗
Minitive measures	✓	✓✓	✓	✗

